



LP187
13/B
White
B-2-23-94

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

To 18X
2-24-94
M21

In re Application of :
: R. BRUCE WALLACE :
: Serial No. 07/996,771 : Group Art Unit 1807
: Filed: December 24, 1992 : Examiner: S. Chambers
: FOR: LIGATION AMPLIFICATION :
: OF NUCLEIC ACID :
: SEQUENCES :

JAN 5 1 1994
GROUP 1800

RESPONSE TO ACTION MAILED JULY 12, 1993

Honorable Commissioner of
Patents and Trademarks
Washington, D. C. 20231

Sir:

Cancel claims 2, 3, 4, 10, 18, 31-35, 37-39, 44 and 45.

Claim 1, lines 1-2, delete "a method for distinguishing
between two known single nucleotide variants"; line 3,
delete "sequence".

Add the following new claims 48, 49 and 50:

--48. An assay for a biologically derived DNA or RNA
test substance, which has a known normal nucleotide sequence
and a known possible mutation at at least one target
nucleotide position in said sequence, which assay determines
whether the test substance has said normal nucleotide
sequence or said possible mutation, said assay comprising
the steps of

- (a) annealing a target oligonucleotide probe of
predetermined sequence to a first sequence of said
test substance so that said target nucleotide
position is aligned with a nucleotide in an end
region of said target probe,

DN20491 01/26/94 07996771 09-0948 020 217 420.00CH

B1
conced

- (b) annealing an adjacent oligonucleotide probe of predetermined sequence to a second sequence of said test substance contiguous to said first sequence, so that the terminal nucleotide in said end region of said target probe and one end of said adjacent probe are directly adjacent to each other,
- (c) contacting said annealed target probe and adjacent probe with a ligase under conditions such that the directly adjacent ends of said probes ligate to form a linked probe product unless there is nucleotide base pair mismatching between said target probe and said test substance at the target nucleotide position,
- (d) separating said test substance and linked probe product, if formed, and
- (e) detecting whether or not said probe product is formed as an indication of nucleotide base pair matching or mismatching at said target nucleotide position.

49. The assay of claim 48 wherein said test substance comprises DNA sequences derived from genomic DNA.

50. The assay of claim 49 wherein said DNA sequences include sequences encoding all or part of normal β -globin or sickle β -globin gene.--

REMARKS

Rejection of Claims 1-4 Under 35 U.S.C. §112, Second Paragraph

Claims 1 to 4 have been amended to avoid this rejection.